

## General Information

So far, so good on the stink bug averages that we are catching. Sheath Blight reached above threshold in the Stoddard (Schuchart) field before reaching the safe zone of 50% heading.

Table 1. Verification field information for the week of July 28th - Aug 3rd.

County	Cultivar	Planted/Rate/Stage	Agronomic Details
Mississippi Sharkey Clay	RT XP753* (FIR)	4/1/24#/ Anthesis or flowering	Stink Bugs average less than 1 per 10 sweeps. 7/29
Stoddard (McMeans) Silty Loam	RT 7521 FP* (FIR)	4/15/22# Anthesis or flowering	Stink Bugs averaged 1 per 10 sweeps. 7/29
Stoddard (Schuchart) Clayey Loam	DG263L+ (FIR)	4/23/40# 10% Heading 7.29	Quadris at 12 oz per acre to suppress sheath blight. 8/1

\*RiceTec Seed Treatment which insecticide, fungicide, zinc, and gibberellic acid.

+Dyna-Gro Seed Treatment which insecticide, fungicide, zinc, and gibberellic acid.

This project would not be possible without the cooperation of the following individuals, Ben Ray, Brandon Copeland, Jennifer and Oliver McMeans, John Davis, Cody Placher, and Paul Schuchart.